

## Exercise Sheet 6

Out: 2018-10-29

Due: 2018-11-05

## Problem 1: Trace Distance

(a) Let  $E_1$  and  $E_2$  be quantum state probability distributions. Let  $\rho_1$  and  $\rho_2$  be the corresponding density operators. Assume that  $E_1$  and  $E_2$  are physically indistinguishable. What is  $\text{TD}(\rho_1, \rho_2)$ ?

(b) Let  $E_1 := \{|+\rangle, \frac{1}{2}\}, \{|-\rangle, \frac{1}{2}\}$  and  $E_2 := \{|0\rangle, 1\}$  be quantum state probability distributions. Let  $\rho_1$  and  $\rho_2$  be the corresponding density operators. What is  $\text{TD}(\rho_1, \rho_2)$ ?

**Note:** You may use computer algebra software (e.g., SageMath) to compute the eigenvectors of matrices if you wish.

(c) Let  $\rho = p\tau + q\rho'$  and  $\sigma = p\tau + q\sigma'$  where  $\tau, \rho', \sigma'$  are density operators, and  $p, q \geq 0$ ,  $p + q = 1$ . Show that  $\text{TD}(\sigma, \rho) = q \cdot \text{TD}(\sigma', \rho')$ .

**Note:** Do not use Lemma 9 in the lecture notes.

(d) Let  $E_1 := \{|+\rangle, \frac{1}{4}\}, \{|-\rangle, \frac{1}{4}\}, (|\Psi\rangle, \frac{1}{2})$ . Let  $E_2 := \{|0\rangle, \frac{1}{2}\}, (|\Psi\rangle, \frac{1}{2})$ . Here  $|\Psi\rangle := \frac{1}{\sqrt{3}}|0\rangle - \sqrt{\frac{2}{3}}|1\rangle$ . Let  $\rho_1$  and  $\rho_2$  be the corresponding density operators. What is  $\text{TD}(\rho_1, \rho_2)$ ?

**Hint:** Consider (c).

(e) Consider the following setup: Alice has a secret bit  $b \in \{0, 1\}$ . Then she chooses randomly  $r \in \{0, 1\}$ . If  $r = 0$ , she encodes  $b$  in the  $|0\rangle, |1\rangle$  basis (i.e., she sends  $|0\rangle$  for  $b = 0$  and  $|1\rangle$  for  $b = 1$ ). If  $r = 1$ , she encodes  $b$  in the  $|+\rangle, |-\rangle$  basis. Then she sends the resulting state  $|\Psi_b\rangle$  to Eve. Show that the trace distance between the mixed states  $\rho_0$  and  $\rho_1$  corresponding to  $b = 0$  and  $b = 1$ , respectively, is  $\text{TD}(\rho_0, \rho_1) = \frac{1}{\sqrt{2}}$ .

**Hint:** The eigenvalues of  $\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{pmatrix}$  are  $\frac{1}{\sqrt{2}}$  and  $-\frac{1}{\sqrt{2}}$ . Note that this is not the toy protocol from the lecture, in the toy protocol  $b$  selected the basis, not  $r$ .

(f) In the experiment described in (e), assume that the bit  $b$  is chosen uniformly at random. Show that Eve cannot guess  $b$  with probability larger than  $\frac{1}{2} + \frac{1}{2\sqrt{2}} \approx 85\%$ .

**Hint:** Try to express the probability that Eve guesses correctly in terms of  $\Pr[G = x|b = y]$  for various  $x, y \in \{0, 1\}$  (here  $G$  denotes Eve's guess) and then use (e).

## Problem 2: Purification (Bonus problem)

Compute purifications of the following density operators:

$$\rho_1 := \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad \rho_2 := \frac{1}{2}|00\rangle\langle 00| + \frac{1}{2}|11\rangle\langle 11|, \quad \rho_3 := \begin{pmatrix} \frac{1}{2} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{2} \end{pmatrix}.$$

**Hint:** Eigenvectors of  $\rho_3$  are  $|+\rangle = \frac{1}{\sqrt{2}}|0\rangle + \frac{1}{\sqrt{2}}|1\rangle$  and  $|-\rangle = \frac{1}{\sqrt{2}}|0\rangle - \frac{1}{\sqrt{2}}|1\rangle$ .